

Finite gap integration of a discrete Euler top

B. Lorbeer*

Fachbereich Mathematik, Technische Universität Berlin, Str.
17 Juni 135, 10623 Berlin, Germany

Abstract

In [1] new discretizations of the Euler top have been found. They can be described with a Lax pair with a spectral parameter on an elliptic curve. This is used in this paper to perform a finite gap integration.

*E-mail: lorbeer@math.tu-berlin.de

1 Introduction

The motion of a rigid body which is not submitted to gravitational force, this is known as the Euler case, is one of the few tops which are completely integrable. These systems have been studied by lots of famous mathematicians like Jacobi and Klein to mention only two.

A new approach to integrable systems in general (finite and infinite dimensional) is provided by the method of finite gap integration which uses Lax pairs with a spectral parameter. This has been introduced by the work of Novikov, Dubrovin, Matveev, Its, Krichever, and others. Once such a Lax pair is found, one can start to use algebro-geometric methods to describe the system. Accounts of what has been achieved up to now can be found in [2].

In the last ten to twenty years completely integrable discrete systems gained more and more attention. Among the early works in this field I only mention [5], [6], [7], [8]. A collection of new articles relevant within this context is [9]. [7] is especially important for the present paper, since it is the first one to give a description of the discrete Euler top with the help of a Lax pair. Here the spectral parameter varies on $\mathbf{CP}^1$. The ideas put down there were elaborated further in [10] and [11].

In [1] we studied another Lax pair for the discrete Euler top where the spectral parameter varies on an elliptic curve. This makes the finite gap integration more difficult. To my knowledge the first solution of such a system was given by A.I. Bobenko in [12]. The present paper can be seen as an addendum to [1]. It describes the finite gap integration for the system introduced there.

2 A Discrete Euler Top

In [1] a discrete version of the Euler top was studied. These notes now are concerned with the finite gap integration of this system. Let's recall the notation:

The top itself is given by the inertia tensor (in diagonal form):

$$\mathbf{I} = \text{diag}(A, B, C), A > B > C > 0. \quad (1)$$

The vectors $\mathbf{M}, \mathbf{\Omega}$ denote the angular momentum and the angular velocity respectively. That is $\mathbf{M} = \mathbf{I}\mathbf{\Omega}$ and the continuous Euler top is given by the well known Euler equation in $\mathbf{R}^3$:

$$\dot{\mathbf{M}} = \mathbf{M} \times \boldsymbol{\Omega}. \quad (2)$$

The discrete version of the Euler top mentioned above is given by a Lax pair depending on a spectral parameter varying on a torus. Let:

$$\begin{aligned} w_1(u) &= \rho \frac{1}{\operatorname{sn}(u, \kappa)}, \\ w_2(u) &= \rho \frac{\operatorname{dn}(u, \kappa)}{\operatorname{sn}(u, \kappa)}, \\ w_3(u) &= \rho \frac{\operatorname{cn}(u, \kappa)}{\operatorname{sn}(u, \kappa)}. \end{aligned} \quad (3)$$

Here sn , cn , dn are the Jacobi elliptic functions and the parameter ρ and the module κ are defined by

$$\begin{aligned} \rho^2 &= \frac{C - A}{AC}, \\ \kappa^2 &= \frac{C(A - B)}{B(A - C)}. \end{aligned} \quad (4)$$

Next let $\mathbf{V} = (V_1, V_2, V_3)^t$ be a vector in $\mathbf{R}^3$, $\sigma_k (k = 1, 2, 3)$ the Pauli matrices and finally let u_0 be such that

$$w_1(u_0) = A^{-\frac{1}{2}}, \quad w_2(u_0) = B^{-\frac{1}{2}}, \quad w_3(u_0) = C^{-\frac{1}{2}}, \quad (5)$$

(such a point exists because of the special choice of the parameters ρ, κ). Now we can define the Lax-pair:

$$\begin{aligned} M &= \frac{1}{2i} \sum_{k=1}^3 M_k w_k(u) \sigma_k, \\ \mathcal{V} &= \mathbf{1} + \frac{h}{2i} \sum_{k=1}^3 V_k w_k(u_0 - u) \sigma_k. \end{aligned} \quad (6)$$

The discrete evolution is given by the Lax-equation

$$M_{n+1}(u) = \mathcal{V}_n(u) M_n(u) \mathcal{V}_n^{-1}(u), \quad (7)$$

which will sometimes also be written as

$$\hat{M}(u) = \mathcal{V}(u)M(u)\mathcal{V}^{-1}(u) . \quad (8)$$

These are the compatibility conditions for the following system:

$$\left| \begin{array}{lcl} M\Phi & = & \Phi\check{\mu} \\ \hat{\Phi} & = & \mathcal{V}\Phi \end{array} \right| \quad (9)$$

Here $\check{\mu} = \text{diag}(\mu_1, \mu_2)$ where μ_1 and μ_2 are the eigenvalues. In [1] it has been shown that (7) is equivalent to a discrete Euler top given by

$$\hat{M} - M = \frac{\gamma h}{4}(M + \hat{M}) \times \mathbf{I}^{-1}(M + \hat{M}). \quad (10)$$

Here h is the discrete parameter and γ a real number which tends to 1 when h tends to zero. See [1] for more considerations on γ (e.g. how γ has to look like so that the discrete map becomes a Poisson map). This paper is concerned with finding explicit solutions for this system with the help of Prym theta functions and abelian differentials. The methods used here are similar to those in [2].

3 The Spectral Curve

To perform the finite gap integration, we first have to get the spectral curve. From (7) it is clear that for each fixed u the eigenvalues of $M(u)$ are independent of n . So the solutions of

$$\det(M(u) - \mu \mathbf{1}) = 0 \quad (11)$$

are constant. Using the identities

$$\begin{aligned} w_2^2 - B^{-1} &= w_1^2 - A^{-1}, \\ w_3^2 - C^{-1} &= w_1^2 - A^{-1}, \end{aligned} \quad (12)$$

(11) is equivalent to

$$\mu^2 = \frac{-1}{4}(M^2 w_1^2 + E - A^{-1} M^2), \quad (13)$$

where $E = \frac{M_1^2}{A} + \frac{M_2^2}{B} + \frac{M_3^2}{C}$.

Let's denote the torus on which the Jacobi elliptic functions are defined by $\mathbf{T}$ (the two generators for the lattice of the torus are given by $4K$ and $4iK'$ where K is the complete elliptic integral of the modulus κ and K' is the complete elliptic integral of the complementary modulus; both are real in this case). Furthermore let

$$\lambda : \mathbf{T} \rightarrow \mathbf{T}, u \rightarrow u + 2K \quad (14)$$

$$\epsilon : \mathbf{T} \rightarrow \mathbf{T}, u \rightarrow u + 2iK'. \quad (15)$$

If u is a point on the torus the set $\{u, \lambda u, \epsilon u, \lambda \epsilon u\}$ will be denoted by $\langle u \rangle$. Next, w_1^2 is invariant w.r.t. λ and ϵ , so the set of solutions of (13) is invariant w.r.t. $(u, \mu) \rightarrow (u + 2K, \mu)$ and $(u, \mu) \rightarrow (u + 2iK', \mu)$ which henceforth will also be denoted by λ and ϵ respectively.

Lemma 1 *The set of solutions of (13) can be made into a Riemann surface.*

Proof: Everywhere except at the points $\langle u_b \rangle$ with $\mu^2(u_b) = 0$ and at the points $\langle 0 \rangle$, u is a chart. At the branch points $\langle u_b \rangle$

$$\mu^2 = [-\frac{M^2}{2}w_1(u_b)w_1'(u_b)](u - u_b) + \dots, u \rightarrow u_b \quad (16)$$

Now let's presume for the moment that $w_1(u_b)w_1'(u_b) \neq 0$. Then μ is obviously a chart.

Next, delete the line $\{(0, z) | z \in \mathbf{C}\}$ from $\mathbf{T} \times \mathbf{C}$, obtaining, say, $(\mathbf{T} \times \mathbf{C})^o$. Let B be a small disk in $\mathbf{C}$ centered at zero, B^o the punctured disk, and glue $B \times \mathbf{C}$ to $(\mathbf{T} \times \mathbf{C})^o$ via the following map m :

$$m : B^o \times \mathbf{C} \rightarrow (\mathbf{T} \times \mathbf{C})^o, (t, \nu) \rightarrow (u, \mu) = (t, \frac{\nu}{t}) \quad (17)$$

In the new coordinates the curve (13) is given by

$$\begin{aligned} \nu^2 &= \frac{-1}{4}t^2(M^2w_1^2(t) + E - A^{-1}M^2) \\ &= \frac{-1}{4}M^2\rho^2 + O(t), t \rightarrow 0. \end{aligned} \quad (18)$$

Thus, adding two points at $(t = 0, \nu = \pm \frac{M^2\rho^2}{4})$, the set of solutions of (13) is compactified to a compact Riemann surface with u a chart at the points $\langle 0 \rangle$. **qed.**

This compact curve is called the spectral curve and will be referred to by the symbol C . It is invariant w.r.t. λ and ϵ . The exchange of sheets will be denoted by π :

$$\pi : C \rightarrow C, (u, \mu) \rightarrow (u, -\mu). \quad (19)$$

For each of the maps λ, ϵ , and π we have natural maps to the belonging quotient spaces, which are all compact Riemann surfaces, too, and these maps will be denoted by $pr_\lambda, pr_\mu, pr_\pi$, respectively. At times, we will also make use of the following map:

$$\chi : C \rightarrow C, (u, \mu) \rightarrow (-u, -\mu). \quad (20)$$

Next, the locations of the branch points have to be described better. They are given by:

$$-w_1^2 + A^{-1} - \frac{E}{M^2} = 0 \quad (21)$$

Using the definition of ρ , κ , and of sn as the inverse of the first elliptic integral, we arrive at the following situation: There are eight branch points, two of them (u_{b1} and u_{b2}) in the small torus $\mathbf{T}/\lambda/\epsilon$ (the others are obtained applying the maps λ, ϵ , and $\lambda\epsilon$). For $\frac{E}{M^2} > \frac{1}{B}$ they are situated on the line $K \rightarrow K + 2iK'$, where $\bar{u}_{b1} = u_{b2}$. For $\frac{E}{M^2} < \frac{1}{B}$ they are situated on the line $iK' \rightarrow 2K + iK'$ and are symmetric over the point $K + iK'$. When $E = \frac{M^2}{C}$ (this corresponds to the stable fix points $(0, 0, \pm|M|)$), the eight branch points degenerate to four points given by $\langle K \rangle$. For $E = \frac{M^2}{A}$ (corresponding to the stable fix points $(\pm|M|, 0, 0)$) we get $\langle iK' \rangle$ and finally for $E = \frac{M^2}{B}$ (corresponding to the separatrix of the unstable fix points $(0, \pm|M|, 0)$) we arrive at $\langle K + iK' \rangle$. Next we find the appropriate cuts. Using the new coordinates (t, ν) at $u = 0$ shows that on a path on C which projects to a straight line through $u = 0$ the function μ changes sign, i.e. starting at (u, μ) we end up at $(u, -\mu)$ (C is invariant w.r.t. the map $(u, \mu) \rightarrow (-u, \mu)$). Thus, going from, say, iK' to $3iK'$ and from K to $3K$, an odd number of cuts should be crossed. Hence an appropriate picture of $C/\lambda/\epsilon$ would be figure (1).




Figure 1: from left to right: $\frac{E}{M^2} > B^{-1}$ and $\frac{E}{M^2} < B^{-1}$

4 The Divisors Of The Baker-Akhiezer Functions

The eigenvectors of the matrices M_n (see (9)) can be understood as functions on the spectral curve: just attach to the point (u, μ) the eigenvector ϕ of $M_n(u)$ with eigenvalue μ (they are, of course, determined only up to multiplication by an arbitrary function on C , so an appropriate normalization will be introduced (see e.g. [3])).

The idea of finite-gap integration is to find as many information as possible about the divisors of the eigenvector functions ϕ_n and then to reconstruct these functions and from them finally the matrices M_n and the vectors $\mathbf{M}_n$. The reason why this is possible is that the considered functions are meromorphic on a *compact* Riemann surface, hence strong tools like the Riemann-Roch theorem can be used.

First we consider the eigenvector function $\phi_0(u, \mu)$ of the matrix $M_{n=0}$ on the spectral curve. It is defined at each point up to a multiplicative number and so it is possible to take the normalized vector function:

$$\phi_0(u, \mu) = \begin{pmatrix} 1 \\ \phi_{0_2} \end{pmatrix}. \quad (22)$$

To find the zeros and poles of ϕ_{0_2} we introduce the following abbreviations:

$$M_0(u) = \begin{pmatrix} \alpha & \beta \\ \delta & -\alpha \end{pmatrix}, \quad (23)$$

$$\mu^2 = \alpha^2 + \beta\delta. \quad (24)$$

Lemma 2 *Let $\check{u} \in \mathbf{T}$ be such that $\beta(\check{u}) = 0$. Then define*

$$P(\check{u}) := (\check{u}, \alpha(\check{u})) + \lambda(\check{u}, \alpha(\check{u})) + \chi(\check{u}, \alpha(\check{u})) + \lambda\chi(\check{u}, \alpha(\check{u})). \quad (25)$$

With this notation the pole divisor of ϕ_{0_2} is given by $P(\check{u})$ and the zero divisor by $\epsilon P(\check{u})$, i.e.:

$$(\phi_{0_2}) = \epsilon P(\check{u}) - P(\check{u}). \quad (26)$$

Proof: We have

$$\phi_{0_2} = \frac{\mu - \alpha}{\beta}, \quad (27)$$

which is a meromorphic function on C . It can only have poles where $\beta = 0$ or $\mu - \alpha = \infty$. Over $\check{u}$ we have $\mu = \pm\alpha$, so there is exactly one pole over $\check{u}$ at $(\check{u}, +\alpha)$ ($\mu = \alpha = 0$ only in the uninteresting case $E = \frac{M^2}{C}$, since $(\alpha)_{\text{zero}} = (\omega_3)_{\text{zero}} = K + \lambda K + \epsilon K + \lambda \epsilon K$.) Now the functions w_k ($k = 1, 2, 3$) have first order poles at the points $\langle 0 \rangle$. Hence β has four poles and thus also four zeros in $\mathbf{T}$. This gives four poles for ϕ_{0_2} in C . Recalling the periodicity of the w_k :

$$\begin{aligned} w_1(\lambda u) &= -w_1(u), & w_1(\epsilon u) &= w_1(u), \\ w_2(\lambda u) &= -w_2(u), & w_2(\epsilon u) &= -w_2(u), \\ w_3(\lambda u) &= w_3(u), & w_3(\epsilon u) &= -w_3(u), \end{aligned} \quad (28)$$

and the fact that all w_k are odd, i.e. $\phi_{0_2}(p) = \phi_{0_2}(\chi p)$, we see that these four poles are given by $P(\check{u})$. The function $\mu - \alpha$ can only have poles over the points $\langle 0 \rangle$ where all w_k have a pole, i.e. β too, so the four poles above are the only ones.

Accordingly, ϕ_{0_2} has four zeros given by $\delta = 0$ (see(24)). From (28) it follows that $\beta(u + 2iK') = \delta(u)$ and $\alpha(u + 2iK') = -\alpha(u)$, thus the zero divisor is given by $\epsilon P(\check{u})$. **qed.**

The next step is the discription of the discrete evolution. We will, for every $n \in \mathbf{N}$, find enough properties of $\phi_n = \mathcal{V}_{n-1} \dots \mathcal{V}_0 \phi_0$, which define this n -th eigenvector function uniquely.

The divisors of ϕ_{n_1}, ϕ_{n_2} are given as follows:

Proposition 3 *Let U be a small neighborhood of u_0 in $\mathbf{T}$ such that $pr_\pi^{-1}(U)$ consists of two nonempty open sets called “the sheets near u_0 ”. Further, let*

ϕ_0 be the initial eigenvector with $(\phi_0)_2 = \epsilon P(\check{u}) - P(\check{u})$ given by the initial data. For h small enough, there exist for each $n \in \mathbf{N}$ a point $\tilde{u}_n$ in $\mathbf{T}$ and two points p_{n_1} and p_{n_2} on C , $p_{n_1}, p_{n_2} \in pr_\pi^{-1}U$, with the following properties:

$$\begin{aligned} - & \Im(pr_\pi p_{n_1}) = \Im(pr_\pi p_{n_2}) = \Im(u_0) \\ - & pr_\pi p_{n_1} = 2u_0 - pr_\pi p_{n_2} \\ - & p_{n_1} \text{ and } p_{n_2} \text{ lie in different sheets,} \end{aligned} \tag{29}$$

from which the divisors of ϕ_{n_1} , ϕ_{n_2} are constructed as follows:

$$(\phi_{n_1}) = \sum_{l=1}^n B_l - P(\check{u}) + P(\tilde{u}_n), \tag{30}$$

$$(\phi_{n_2}) = \sum_{l=1}^n B_l - P(\check{u}) + \epsilon P(\tilde{u}_n), \tag{31}$$

$$B_n := E_n + \lambda E_n + \epsilon E_n + \lambda \epsilon E_n, \tag{32}$$

$$E_n := -(u_0, \mu) - (u_0, -\mu) + p_{n_1} + p_{n_2}. \tag{33}$$

If h tends to zero, we get $\lim_{h \rightarrow 0} pr_\pi p_{n_k} = u_0$ ($k = 1, 2$). See figure (2).

Conversely, for each given set of divisors $\{B_n | n \in \mathbf{N}, n > 0\}$, of the form (32) with $pr_\pi p_{n_1} - u_0$ small, there is a sequence of vector functions ϕ_n with divisors (30), (31), which satisfy (9).

Proof: First we note that multiplication with $\mathcal{V}_0$ adds eight poles to each component of ϕ_0 , one pole at each point over each element of $\langle u_0 \rangle$. We prove this for the coefficient ϕ_{0_1} , for the second the procedure is similar.

Recall that

$$\frac{\pm \mu - \alpha}{\beta}(u_0) = \frac{\pm \frac{i}{2} \sqrt{E} - \frac{1}{2i} \frac{M_3}{\sqrt{C}}}{\frac{1}{2i} (\frac{M_1}{\sqrt{A}}) - i \frac{M_2}{\sqrt{B}}}, \tag{34}$$

and

$$V_k w_k(u_0 - u) = (\rho V_k)(u_0 - u)^{-1} + O(1), \quad u \rightarrow u_0, \quad (k = 1, 2, 3), \tag{35}$$

and for $h \rightarrow 0$ we have $\mathbf{V} \rightarrow (\frac{M_1}{\sqrt{A}}, \frac{M_2}{\sqrt{B}}, \frac{M_3}{\sqrt{C}})^t$ (see [1]).

Hence, if no new poles appear, we must have:




Figure 2: possible divisor of ϕ_{21} (to get the divisor of ϕ_{22} use ϵ on the four “separate” (away from $\langle u_0 \rangle$) zeros

$$\begin{aligned} \frac{M_3}{\sqrt{C}} + \left(\frac{M_1}{\sqrt{A}} - i \frac{M_2}{\sqrt{B}} \right) \frac{\pm \frac{i}{2} \sqrt{E} - \frac{1}{2i} \frac{M_3}{\sqrt{C}}}{\frac{1}{2i} \left(\frac{M_1}{\sqrt{A}} \right) - i \frac{M_2}{\sqrt{B}}} &= 0 \\ \iff \sqrt{E} &= 0, \end{aligned}$$

which happens only in the uninteresting case $\mathbf{M} = 0$.

Thus each component of ϕ_1 has 4+8 poles: four poles given by the initial data (the poles of ϕ_{02}) and the eight poles just considered. But we know that the divisor of $\frac{\phi_{12}}{\phi_{11}}$ is given by

$$\left(\frac{\phi_{12}}{\phi_{11}} \right) = \epsilon P(\tilde{u}_1) - P(\tilde{u}_1) \quad (36)$$

for some $\tilde{u}_1$ (the same proof as for ϕ_{02} , see above). Whence, since *all* poles coincide, eight of the 12 zeros of ϕ_{11} coincide with eight zeros of ϕ_{12} . At those points $\det \mathcal{V}_0$ must be zero.

Now:

$$\begin{aligned} \det \mathcal{V} &= 0 \\ \iff w_1^2(u_0 - u) &= \mathbf{V}^{-2} \left(\frac{-4}{h^2} + A^{-1} \mathbf{V}^2 - \langle \mathbf{V}, \mathbf{I}^{-1} \mathbf{V} \rangle \right). \end{aligned} \quad (37)$$

$w_1^2(u_0 - u)$ is a well defined function on $\mathbf{T}/\lambda/\epsilon$ and having there only one pole of order two, it has also two points u_{1_1}, u_{1_2} where (37) is satisfied, thus in $\mathbf{T}$ (37) is true at the points $\langle u_{1_1} \rangle \cup \langle u_{1_2} \rangle$, i.e. there are 16 possible sites for the eight zeros of ϕ_1 . Since $\mathbf{V}$ is a real vector and for small h the right hand side of (37) converges to $-\infty$, we get (recall the definition of w_1):

$$\Im(u_{1_1}) = \Im(u_{1_2}) = \Im(u_0), \quad (38)$$

and since w_1 is odd:

$$u_{1_1} + u_{1_2} = 2u_0, \quad (39)$$

and for $h \rightarrow 0 : u_{1_1}, u_{1_2} \rightarrow u_0$. Next, $\mathcal{V}_0 \neq 0$ for all u , which means that there is exactly one zero over each element of $\langle u_{1_1} \rangle \cup \langle u_{1_2} \rangle$. These zeros are invariant w.r.t. λ and ϵ :

First note that:

$$\begin{aligned} M(\lambda u) &= \sigma_3 M(u) \sigma_3 \Rightarrow \phi_0(\lambda p) = \sigma_3 \phi_0(p), \\ M(\epsilon u) &= \sigma_1 M(u) \sigma_1 \Rightarrow \phi_0(\epsilon p) = \phi_{0_2}^{-1}(p) \sigma_1 \phi_0(p), \\ \mathcal{V}(\lambda u) &= \sigma_3 \mathcal{V}(u) \sigma_3, \\ \mathcal{V}(\epsilon u) &= \sigma_1 \mathcal{V}(u) \sigma_1. \end{aligned} \quad (40)$$

Thus, if

$$\mathcal{V}_0(pr_\pi p) \phi_0(p) = 0, \quad (41)$$

then

$$\mathcal{V}_0(\lambda pr_\pi p) \phi_0(\lambda p) = \mathcal{V}_0(\epsilon pr_\pi p) \phi_0(\epsilon p) = 0. \quad (42)$$

Let p_{1_1}, p_{1_2} be the two zeros with $pr_\pi p_{1_i} = u_{1_i}$, ($i = 1, 2$). We now show that p_{1_1} and p_{1_2} lie in different sheets: Let $W_0(u) := \mathcal{V}_0(u) - \mathbf{1}$, then $\mathcal{V}_0(u_{1_2}) = \mathcal{V}_0(u_{1_1}) - 2W_0(u_{1_1})$, since all w_k are odd. Next, let $\mathcal{V}_0(u_{1_1}) \phi_0(u_{1_1}, \mu) = 0$. Then:

$$\begin{aligned}
\mathcal{V}_0(u_{12})\phi_0(u_{12}, \mu) &= (\mathcal{V}_0(u_{11}) - 2W_0(u_{11}))(\phi_0(u_{11}, \mu) + O(h)) \\
&= -2W_0(u_{11})\phi_0(u_{11}, \mu) + O(h) \\
&= 2\phi_0(u_{11}, \mu) + O(h), \quad h \rightarrow 0,
\end{aligned} \tag{43}$$

using the definition of $W_0(u)$. Since $\phi_0(u_{11}, \mu)$ doesn't vanish, the second zero of ϕ_1 must lie on the other sheet.

Exchanging the sheets for p_{11} and p_{12} obviously means changing the sign of h . It is clear that further multiplications with the matrices $\mathcal{V}_n(u)$ lead to the increase of the order of the poles over $\langle u_0 \rangle$ and to further pairs of zeros p_{n1}, p_{n2} common to ϕ_{n1} and ϕ_{n2} near the points $\langle u_0 \rangle$. The proof that $(\frac{\phi_{n2}}{\phi_{n1}}) = \epsilon P(\tilde{u}_n) - P(\tilde{u}_n)$ for some $\tilde{u}_n$ is the same as for ϕ_{02} .

Conversely, given a set of divisors $\{B_n | n \in \mathbf{N}, n > 0\}$ with the above properties we only have to vary $h\gamma$ for each step to arrive at a sequence of angular momentum vectors $\mathbf{M}_n$ which belong to eigenvector functions with divisors given by the prescribed set of B_n divisors. **qed.**

Below we will also need the following properties:

$$\phi_n(\lambda p) = \sigma_3 \phi_n(p), \tag{44}$$

since $(\mathcal{V}_{n-1} \cdots \mathcal{V}_0 \phi_0)(\lambda u) = (\sigma_3 \mathcal{V}_{n-1} \sigma_3 \cdots \sigma_3 \mathcal{V}_0 \sigma_3 \phi_0)(p) = \sigma_3 \phi_0(p)$. Similarly:

$$\phi_n(\epsilon p) = \phi_{02}^{-1}(p) \sigma_1 \phi_n(p). \tag{45}$$

Also recall that the four “separate” zeros of ϕ_{n1} and ϕ_{n2} have the symmetry χ :

$$\chi P(\tilde{u}_n) = P(\tilde{u}_n). \tag{46}$$

Proposition 4 *If ψ_n and ϕ_n are two holomorphic vector functions on the spectral curve which both have divisors as in proposition (3) and satisfy (44), (45), and (46), then $\phi_n = c\psi_n$, $c \in \mathbf{C}^*$.*

Proof: Let ϕ_{n1}, ψ_{n1} be the first coefficients of the vector functions. Then $\frac{\phi_{n1}}{\psi_{n1}}$ is a function on the torus C/λ with two poles at $D = p + \chi p$ for some

p . Two of the three holomorphic differentials on this compact curve are du and $\frac{du}{\mu}$ (here u is the pullback of the chart u from $\mathbf{T}$ to C). Its values in the chart u at the poles are given by:

$$\begin{array}{c|cc} & du & \frac{du}{\mu} \\ \hline p & 1 & \mu^{-1} \\ \chi p & 1 & -\mu^{-1} \end{array} \quad (47)$$

for some $\mu \neq 0$. The rank of this matrix is two, thus the index of speciality of D is 1. From the Riemann-Roch formula it follows that the space of holomorphic functions f with $(f) + D$ being integral is one. But the constant functions satisfy this condition, so they are the only ones. Hence $\frac{\phi_{n_1}}{\psi_{n_1}} = c = \text{const.}$

But from (45) we see that also $\frac{\phi_{n_2}}{\psi_{n_2}} = c = \text{const.}$ (the same constant c as for the first coefficients). Thus $\phi_n = c\psi_n$. **qed.**

5 The Prym Theta Function

In this section the construction of a Prym theta function is described. This will then be used in the formulas for ϕ_n .

Consider the Riemann surface C/λ . We will first investigate the case $\frac{E}{M^2} > B^{-1}$. The homology basis is chosen as in figure (3). That is

$$\begin{aligned} \pi a_1 &= -a_3, & \pi b_1 &= -b_3, \\ \pi a_2 &= -a_2, & \pi b_2 &= -b_2, \end{aligned} \quad (48)$$

The corresponding normalized cohomology basis will be denoted by v_k , ($k = 1, 2, 3$):

$$\int_{a_k} v_l = 2\pi i \delta_{kl}. \quad (49)$$

A Prym differential for the projection π is given by

$$\omega := v_1 + v_3. \quad (50)$$

We then have

$$\pi^* \omega = \pi^*(v_1 + v_3) = -v_3 - v_1 = -\omega \quad (51)$$




Figure 3: appropriate homology basis for $\frac{E}{M^2} > B^{-1}$

(For a thorough discussion of Prym differentials and Prym varieties see [4]. In our particular case the differential ω can be given explicitly by $\omega = \frac{2\pi i}{c} \frac{du}{\mu}$ where $c := \int_{a_1} \frac{du}{\mu}$, but this fact won't be used.)

Lemma 5 *We have*

$$\int_{a_1} \omega = \int_{a_3} \omega = 2\pi i, \int_{a_2} \omega = 0, \quad (52)$$

$$\int_{b_1} \omega = \int_{b_3} \omega =: \tau, \quad (53)$$

$$\int_{b_2} \omega = 2\pi i. \quad (54)$$

Proof: The assertions (52) are obvious from the definition of ω . Further:

$$\int_{b_3} \omega = \int_{-\pi b_1} \omega = - \int_{b_1} \pi^* \omega = \int_{b_1} \omega. \quad (55)$$

Finally, to prove (54), first note that




Figure 4: in the homology group: $b_2 + \epsilon b_2 = a_1 + a_3$

$$b_2 + \epsilon b_2 = a_1 + a_3, \quad (56)$$

(see figure (4)).

Moreover

$$\epsilon^* \omega = \omega. \quad (57)$$

This is true, since the a -periods of $\epsilon \omega$ and ω coincide:

$$\epsilon a_1 = a_1, \quad (58)$$

$$\epsilon a_3 = a_3, \quad (59)$$

so the a_1 - and a_3 -periods coincide. Because of

$$a_2 + \epsilon a_2 = 0, \quad (60)$$

(see figure (5)),

we also have:




Figure 5: In the homology group $a_2 + \epsilon a_2 = 0$

$$\int_{a_2} \epsilon^* \omega = - \int_{\epsilon a_2} \epsilon^* \omega = - \int_{a_2} \omega = 0, \quad (61)$$

so indeed $\epsilon^* \omega = \omega$. Finally, using (56) and (57) it follows

$$\int_{b_2} \omega = \frac{1}{2} \left(\int_{b_2} \omega + \int_{\epsilon b_2} \epsilon^* \omega \right) = \frac{1}{2} \int_{a_1 + a_3} \omega = 2\pi i. \quad (62)$$

qed.

Next, the torus given by the two generators $(2\pi i, \tau)$ will be denoted by J . Let p_0 be an arbitrary point on C/λ , then

$$A(p) := \int_{p_0}^p \omega \quad (63)$$

is a well defined map from C/λ to J because of lemma (5). The theta function on J is defined by:

$$\theta(z, \tau) := \sum_{m \in \mathbf{Z}} \exp\left(\frac{1}{2} m^2 \tau + mz\right). \quad (64)$$

The periods of $\theta(z, \tau)$ are given by ($e \in J$, arbitrary):

$$\theta(z + e + 2\pi i) = \theta(z + e) \quad (65)$$

$$\theta(z + e + \tau) = \exp\left(-\frac{\tau}{2} - z - e\right)\theta(z + e), \quad (66)$$

from which the periods of the pullback $\theta(A(p) + e)$ of $\theta(z + e)$ to C/λ follow.

Proposition 6 *The function $\theta(A(p) + e)$ has two zeros in C/λ and if $\check{p}$ is one of them, then $\chi\check{p}$ is the other.*

Proof: The proof that $\theta(A(p) + e)$ has two zeros in C/λ follows the standard procedure: just cut C along the homology basis and then integrate $\frac{d\theta(z+e)}{\theta(z+e)}$ along the image of that cut in J .

To prove the second part of the proposition, first note, that $\chi^*\omega = \omega$: this follows from:

$$\begin{aligned} \chi a_1 &= a_3, \\ \chi a_2 &= -a_2. \end{aligned} \quad (67)$$

So let's choose p_0 to be one of the points over $u = 0$. Let $\check{p}$ be a zero of $\theta(A(p) + e)$ and ζ a path from p_0 to $\check{p}$, i.e. $\chi\zeta$ is a path from p_0 to $\chi\check{p}$. Then:

$$A(\check{p}) = \int_{\zeta} \omega = \int_{\chi\zeta} \chi^*\omega = \int_{\chi\zeta} \omega = A(\chi\check{p}), \quad (68)$$

which proves the second part. **qed.**

For the case $\frac{E}{M^2} < B^{-1}$ we choose the homology basis given in figure (6). With this choice, everything works as in the other case above.

6 The Baker-Akhiezer Function

Now we formulate the main theorem. Let's recall the notations:

Let $\mathbf{M}_0 \in \mathbf{R}^3$ be an initial value for the discrete Euler top, C the spectral curve defined by $\mathbf{M}_0$ and let $P(\check{u})$ be the pole divisor of ϕ_{0_2} (ϕ_0 is the eigenvector function of M_0 with $\phi_{0_1} = 1$). ω and θ will still denote the Prym differential and theta function described above, and




Figure 6: appropriate homology basis for $\frac{E}{M^2} < B^{-1}$

$$e := - \left(\int^{(\check{u}, \alpha(\check{u}))} \omega \right) + \pi i + \frac{\tau}{2}. \quad (69)$$

Further, to each $n \in \mathbf{N}$ we associate the divisors B_n, E_n as in (32) and (33). On the curve $C/\lambda/\epsilon$ let $d\Omega_n$ be the third kind abelian differential with zero a -periods (the a -cycles on $C/\lambda/\epsilon$ (let's denote them by $\tilde{a}$) are defined by:

$$2\tilde{a}_1 = pr_\epsilon a_1 \quad (70)$$

$$2\tilde{a}_2 = pr_\epsilon a_3 \quad (71)$$

and only single poles with residues given by the divisor E_n

$$res_p d\Omega = E_n(p), \quad p \in C/\lambda/\epsilon. \quad (72)$$

Finally,

$$V_n = \int_{b_1} d\Omega_n. \quad (73)$$

In what follows, pullbacks of differentials are usually denoted with the same symbol as the original differentials and it will be clear from the context which one is meant.

Theorem 7 *The functions:*

$$\phi_{n_1}(p) := \frac{\theta(\int^p \omega + \sum_{k=1}^n V_k + e)}{\theta(\int^p \omega + e)} \exp\left(\int^p \sum_{k=1}^n d\Omega_k\right) \quad (74)$$

$$\phi_{n_2}(p) := \frac{\theta(\int^p \omega + \sum_{k=1}^n V_k + \pi i + e)}{\theta(\int^p \omega + e)} \exp\left(\int^p \sum_{k=1}^n d\Omega_k\right) \quad (75)$$

are well defined on C and are the coefficient functions of an eigenvector function of M_n .

Proof: We first prove that ϕ_{n_1} and ϕ_{n_2} are functions on C . Let's consider C/λ . The a -cycles cause no problems. For the b_2 -cycle, note that

$$\int_{b_2} d\Omega_k = \frac{1}{2} \left(\int_{b_2} d\Omega_k + \int_{\epsilon b_2} \epsilon^* d\Omega_k \right) = \frac{1}{2} \int_{a_1+a_3} d\Omega_k = 0. \quad (76)$$

Next we show that $\int_{b_1} d\Omega_k = \int_{b_3} d\Omega_k$. Since $d\Omega_k$ is a normalized abelian differential of the third kind, we have the Riemann-Roch relations:

$$\int_{b_1} d\Omega_k = \int_{E_n + \epsilon E_n} v_1, \quad (77)$$

$$\int_{b_3} d\Omega_k = \int_{E_n + \epsilon E_n} v_3, \quad (78)$$

Now $v_1 - v_3 = \frac{2\pi i}{4iK'} du$, since they obviously have the same a -periods. But

$$\int_{E_n + \epsilon E_n} du = 0 \quad (79)$$

and thus the b_1 - and b_3 -periods of $d\Omega_k$ are the same. Hence ϕ_{n_1} does not change w.r.t. any cycle in C/λ while ϕ_{n_2} changes sign when going around b_1 and b_3 in C/λ , i.e. on C it is single valued too.

Now we proceed to show that $\phi_n = (\phi_{n_1}, \phi_{n_2})^t$ is indeed an eigenvector function of M_n . Recall that we have to show the following facts:

- (i) both ϕ_{n_1} and ϕ_{n_2} have four poles given by $P(\check{u})$ (determined by the initial value),

- (ii) both ϕ_{n_1} and ϕ_{n_2} have poles and zeros given by the divisor $\sum_{k=1}^n B_k$,
- (iii) the remaining four zeros in the divisors of ϕ_{n_1} and ϕ_{n_2} are not the same but in both cases they are symmetric w.r.t. the involution χ ,
- (iv) $\phi_n(u + 2K, \mu) = \sigma_3 \phi_n(u, \mu)$,
- (v) $\phi_n(u + 2iK', \mu) = \phi_{0_2}^{-1}(u, \mu) \sigma_1 \phi_n(u, \mu)$.

The fact (i) follows from the appropriate choice of the point e in the torus J . ϕ_{n_1} and ϕ_{n_2} thus both have a pole at $(\check{u}, \alpha(\check{u}))$. From proposition 6 and the behaviour of θ when shifted by τ , we find that $\theta(A(p) + e)$ has zeros at $P(\check{u})$.

Point (ii) is provided by the exponential terms.

Next, we know already that the second coefficient of an eigenvector function of a matrix M of the form $\phi = (1, \phi_2)^t$ has a divisor symmetric w.r.t. the involution χ . Now the pole and zero divisors of $\phi_n = (1, \phi_{n_2})^t$ are given by the four zeros of ϕ_{n_1} and ϕ_{n_2} respectively, which proves (iii).

(iv) follows immediately from the periodic behaviour of θ .

Finally, since ω as well as $d\Omega_k$ are pullbacks via the quotient map $C/\lambda \rightarrow C/\lambda/\epsilon$, and a_1 - and a_3 -periods for ω are $2\pi i$ and for $d\Omega_k$ zero, we get:

$$\begin{aligned} \phi_{n_1}(u + 2iK', \mu) &= \\ &= \frac{\theta(\int^{(u, \mu)} \omega + \sum V_k + \pi i + e)}{\theta(\int^{(u, \mu)} \omega + \pi i + e)} \exp(\int^{(u, \mu)} \sum d\Omega_k), \end{aligned} \quad (80)$$

$$\begin{aligned} \phi_{n_2}(u + 2iK', \mu) &= \\ &= \frac{\theta(\int^{(u, \mu)} \omega + \sum V_k + 2\pi i + e)}{\theta(\int^{(u, \mu)} \omega + \pi i + e)} \exp(\int^{(u, \mu)} \sum d\Omega_k). \end{aligned} \quad (81)$$

Hence:

$$\begin{aligned} \phi_n(u + 2iK', \mu) &= \frac{\theta(\int^{(u, \mu)} \omega + e)}{\theta(\int^{(u, \mu)} \omega + \pi i + e)} \sigma_1 \phi_n(u, \mu) \\ &= \phi_{0_2}^{-1}(u, \mu) \sigma_1 \phi_n(u, \mu), \end{aligned} \quad (82)$$

which proves (v).

qed.

7 Solutions For The Angular Momentum

To get $\mathbf{M}$ from the Baker-Akhiezer functions, we consider (9) at the point $u = 0$. Here the asymptotics of M_n and μ are given by

$$M_n = \left(\frac{\rho}{2i} \sum M_k \sigma_k\right) u^{-1} + O(1), \quad u \rightarrow 0, \quad (83)$$

$$\mu = \pm \left(\frac{\rho}{2i} |M|\right) u^{-1} + O(1), \quad u \rightarrow 0, \quad (84)$$

Using as base p_0 for the integrals a point with $pr_\pi p_0 = 0$, we see that the two eigenvectors at $u = 0$ can, after multiplication with appropriate factors, be written as

$$\Phi_n(p_0) = \begin{pmatrix} 1 & 1 \\ \frac{\theta(\pi i + \sum V_k + e)}{\theta(\sum V_k + e)} & \frac{\theta(d + \pi i + \sum V_k + e)}{\theta(d + \sum V_k + e)} \end{pmatrix} =: \begin{pmatrix} 1 & 1 \\ A & B \end{pmatrix}, \quad (85)$$

where $d := \int_{p_0}^{\pi p_0} \omega$. Using this, (9) leads to

$$\begin{pmatrix} M_3 + (M_1 - iM_2)A & M_3 + (M_1 - iM_2)B \\ M_1 + iM_2 - M_3A & M_1 + iM_2 - M_3B \end{pmatrix} = \begin{pmatrix} M & -M \\ AM & -BM \end{pmatrix}. \quad (86)$$

This finally yields

$$\begin{aligned} M_1 &= M \frac{\theta(\sum V_n + e)\theta(d + \sum V_n + e) - \theta(\pi i + \sum V_n + e)\theta(d + \pi i + \sum V_n + e)}{\theta(\pi i + \sum V_n + e)\theta(d + \sum V_n + e) - \theta(\sum V_n + e)\theta(d + \pi i + \sum V_n + e)} \\ M_2 &= iM \frac{\theta(\pi i + \sum V_n + e)\theta(d + \pi i + \sum V_n + e) + \theta(\sum V_n + e)\theta(d + \sum V_n + e)}{\theta(\pi i + \sum V_n + e)\theta(d + \sum V_n + e) - \theta(\sum V_n + e)\theta(d + \pi i + \sum V_n + e)} \\ M_3 &= M \frac{\theta(\pi i + \sum V_n + e)\theta(d + \sum V_n + e) + \theta(\sum V_n + e)\theta(d + \pi i + \sum V_n + e)}{\theta(\sum V_n + e)\theta(d + \pi i + \sum V_n + e) - \theta(\pi i + \sum V_n + e)\theta(d + \sum V_n + e)}. \end{aligned} \quad (87)$$

8 Acknowledgements

I want to thank A.Bobenko, R.Seiler, and J.Suris for lots of helpful discussions.

9 Bibliography

- [1] A.I.Bobenko, B.Lorbeer, Y.Suris. Integrable discretizations of the Euler top, Preprint No. 312, SFB 288 "Differentialgeometrie und Quantenphysik"
- [2] E.D.Belokolos, A.I.Bobenko, V.Z. Enol'ski, A.R. Its, V.B. Matveev. Algebro-geometric approach to nonlinear integrable equations, Springer, Berlin (1994)
- [3] B. Dubrovin, I. Krichever, S.Novikov. Integrable Systems I, In: Dynamical systems IV. Encyclopaedia of mathematical sciences (vol.4), V.I. Arnol'd, S.P. Novikov (Eds.)
- [4] J.Fay. Theta functions on Riemann surfaces, Lecture Notes in Mathematics 352, Springer, Berlin (1973)
- [5] M.Ablowitz, J. Ladik. A nonlinear difference scheme and inverse scattering. Stud. Appl. Math. 55 (1976) 213-229; On the solution of a class of nonlinear partial difference equations. Stud. Appl. Math. 57 (1977) 1-12.
- [6] F.W. Nijhoff, H.Capel, and G.Wiersma. Integrable lattice systems in two and three dimensions. Lect. Notes Phys. 239 (1984) 263-302; Linearizing integral transform for the multicomponent lattice KP. Physica A 138 (1986) 76-99.
- [7] A.P. Veselov. Integrable systems with discrete time and difference operators. Funct.Anal. Appl. 22 (1988) 1-13.
- [8] Yu.B. Suris. Generalized Toda Chains in discrete time. Algebra i Anal. 2 (1990) 141-157; Discrete-time generalized Toda lattices: complete integrability and relation with relativistic Toda lattices. Phys. Lett. A 145 (1990) 113-119.
- [9] Discrete Integrable Geometry and Physics, Eds. A.Bobenko and R.Seiler, Oxford University Press, 1998.
- [10] J.Moser, A.P. Veselov. Discrete Versions of some classical integrable systems and factorization of matrix polynomials. Commun. Math. Phys. 139 (1991) 217-243.
- [11] P.Deift, L.-Ch.Li, and C.Tomei. Loop groups, discrete versions of some classical integrable systems, and rank 2 extensions. Mem. Amer. Math. Soc. 479 (1992).
- [12] A.I. Bobenko. Real algebro-geometric solutions of the Landau-Lifshits equation in Prym theta functions. Functional Anal. Appl. 19, 5-17 (1985).